


ThinkAskLearn
Health Professional Education

Assessing Suicide Risk in Kids

David Corkill
Emergency Nurse Educator
MEmergN, MAdvPrac (HthProfEdu), BN, Dip App Sc


www.thinkasklearn.com.au



1

Beware Confronting Issues Ahead

- Need help
 - Lifeline 13 11 14
 - <http://www.lifeline.org.au>
 - Beyond Blue
 - www.beyondblue.org.au
 - Kids Helpline
 - 1800 55 1800



2



MENTAL HEALTH FOR EMERGENCY DEPARTMENTS

A REFERENCE GUIDE







3


4




5




6




7



8



9



10

What do the numbers tell us

- Increasing number of suicides, rates steady over 10 years
- Young people are of a concern
- 40-50 years are highest risk
- Another risk group is over 85 years
- Suicide rates – different definition per state
 - Data skewed in youth suicide – underreported?
- But they also only tell us those that have died

11

Risk Factors for Suicide – Demographic

- Male
 - Remember death rates
 - No difference between sexes (my belief/experience)
- Older adolescence (versus younger)
- Non-heterosexual orientation

12

Risk Factors for Suicide – Clinical

- Psychiatric diagnosis
- Recent discharge from psychiatric hospital
- Past suicide attempt
- Family history of suicide
- Child sexual abuse/rape
- Childhood history of trauma



13

Risk Factors for Suicide – Family/Environment

- Life stresses,
 - unemployment and legal and school problems
- Access to lethal means
- Lack of social supports
- Contagion – exposure to others demonstrating suicidal behaviour (imitation, coping strategy)
- Non-intact families (eg. Divorce)



14

Assess the Risk of Suicide

- Assess the risk profile



15

Assess the Risk of Suicide

- Assess the risk profile
- Use open ended questions



16

Ask about suicide



17

Nice Quote

“It should be noted that asking an adolescent if they have suicidal ideation does not make them suicidal; rather, inquiry about suicidal thinking is likely to lead the adolescent to feel they are being listened to and that they are not alone.”

Gordon & Melville 2014



18

Open Ended Questions

- Do you ever feel like giving up?
- Do your symptoms/things ever become too much to cope with?
- Do you ever feel hopeless about your situation?



19

Closed Ended Questions

- Follow open with closed questions
- Do you ever think about going to sleep and not waking up?
- Do you think that you would be better off if you weren't alive?



20

Suicide Risk Assessment

- Thoughts of Suicide but no formal plan
- Plan of action but has no undertaking
- Acted out plan but now regrets attempt
- Acted out plan but no regrets attempt



21

Assess the Risk of Suicide

- Assess the risk profile
- Use open ended questions
- Consider involving the parents if at all possible
- Document the risk assessment, mental state, safety plan and people contacted
- Arrange for review



22

Ways to Show a Suicidal Person You Care

- ☐ If the person is high risk, do not leave them alone
- ☐ Let the person know you are deeply concerned
- ☐ Listen attentively to everything
- ☐ Do not keep suicidal tendencies a secret
- ☐ Don't be judgmental
- ☐ Be careful of the statement that you make
- ☐ Follow up on a regular basis
- ☐ Comfort the person with words of encouragement

Source: suicide.org

23

Jane is 27 years of age and has a depressive illness. She has fluctuating mood symptoms and suicidal thoughts. The urge to follow through with these thoughts generally lasts for 1-2 hours, but then subsides. Jane has reported an awareness that the urge to harm herself will pass if she is able to use her safety plan.

Warning signs/symptoms suggesting that Jane is deteriorating:

- sleeping 7 hours per night
- losing weight
- not going to school
- having suicidal thoughts

Internal coping strategies (things that Jane can do to soothe herself):

- Jane has agreed to make a list of reasons for living and can look at this list if she is struggling
- Jane has also agreed to make a flipbook, which includes pictures of family holidays and her Year 11 formal. Jane can also look at these pictures if she is struggling

Distraction techniques:

- Listen to music, play piano, guitar
- Watch family DVD's
- Have a bath
- Play with her dog, Storage
- Go for a walk

Jane can ask her mother for extra (pres) medication of diazepam 2 mg if she is very distressed and feels that she is not coping.

Social supports that can help take Jane's mind off problems:

- Phone or text Julie (sister)
- Phone or text Julie (sister)

Family and friends for crisis help:

Jane has identified 5 people she can call if she is not able to cope or keep herself safe.

Jane agrees to keep a fully charged mobile telephone and have credit on her phone.

The 5 people who agree to be called on an emergency people are:

1. Mum _____ mobile number _____
2. Dad _____ mobile number _____
3. Susan (friend) _____ mobile number _____
4. Kim (friend) _____ mobile number _____
5. Beverly (sister) _____ mobile number _____

Professionals and agencies:

- Jane can call R2248 helpline 1800 95 1800.
- Jane's mother will call Dr Jones if these strategies do not work and Jane is not coping. Dr Jones can be called during the day on 9123 4567 or after hours on 0204 0886, pager 2345.
- In a crisis, Jane's mother should take her to the Emergency Department at St. Andrews Medical Centre.



24

Jane is 17 years of age and has a depressive illness. She has fluctuating mood symptoms and suicidal thoughts. The urge to follow through with these thoughts generally lasts for 1–2 hours, but then subsides. Jane has reported an awareness that the urge to harm herself will pass if she is able to use her safety plan.



25

Distraction techniques:

- Listen to music, play piano, guitar
- Watch family DVDs
- Have a bath
- Play with her dog, Savage
- Go for a walk

Jane can ask her mother for extra (prn) medication of diazepam 2 mg if she is very distressed and feels that she is not coping.



26

Family and friends for crisis help:

Jane has identified 5 people she can call if she is not able to cope or keep herself safe. Jane agrees to keep a fully charged mobile telephone and have credit on her phone.

The 5 people who agree to be called on as safety people are:

1. Mum _____ mobile number _____
2. Dad _____ mobile number _____
3. Jessie (friend) _____ mobile number _____
4. Kim (friend) _____ mobile number _____
5. Bevetley (aunt) _____ mobile number _____



27

A Quick Case

- A 14yr old female presents to the clinic with a very anxious friend
- She looks pale, shaky and frightened
- Is able to answer questions
- C/o Dizziness,
- *"My face is tingling and my hands are cold? Am I going to die?"*



28

A Quick Case

- Vital signs normal
- SpO₂ 100% Room Air
- C/O Pain with the BP cuff
- Hand cramping
- Anxious
- What do you think is going on?



29

What is wrong with the patient?

- | | |
|---|---|
| • Acute Respiratory Distress Syndrome | • Panic Disorders |
| • Asthma | • Pleural Effusion |
| • Atrial Fibrillation | • Pneumonia, Bacterial |
| • Atrial Flutter | • Pneumothorax, Iatrogenic, Spontaneous and Pneumomediastinum |
| • Cardiomyopathy, Dilated | • Pneumothorax, Tension and Traumatic |
| • Cardiomyopathy, Restrictive | • Pulmonary Embolism |
| • Chronic Obstructive Pulmonary Disease and Emphysema | • Respiratory Distress Syndrome, Adult |
| • Costochondritis | • Smoke Inhalation |
| • Diabetic Ketoacidosis | • Toxicity, Carbon Monoxide |
| • Hyperventilation Syndrome | • Toxicity, Carbon Monoxide |
| • Metabolic Acidosis | • Venous Air Embolism |
| • Methemoglobinemia | • Withdrawal Syndromes |
| • Myocardial Infarction | |
| • Nasopharyngeal Stenosis | |

Kern et al 2014



30

Hyperventilation Syndrome

- Agitation,
- Deep breathing and tachypnea,
- Chest pain, dyspnea,
- Wheezing, dizziness, palpitations,
- Tetanic cramps (eg, carpopedal spasm), paresthesias,
- Generalized weakness,
- "A sense of suffocation"
- An emotionally stressful precipitating event can often be identified.



31

ABNORMAL OXYGEN SATS

ARE NOT CONSISTENT

WITH HYPERVENTILATION SYNDROME

32

NORMAL OXYGEN SATS

DO NOT EXCLUDE

OTHER RESPIRATORY CAUSES

33

What is really happening?

- Minute ventilation exceeds metabolic demands
- Haemodynamic and chemical changes that produce characteristic symptoms
- Drop in arterial partial pressure of carbon dioxide (PaCO_2) through voluntary hyperventilation
- But many patients do not have a low PaCO_2 !!!!



34



35

Ban the Bag – What?

- Death can (have) occurred
 - Misdiagnosis – MI, PE, PNx
- It will not help some patients (CO_2 not low)
- Often unsuccessful in reversing hyperventilation
- Simple reassurance with explanation of how the symptoms may be occurring



36



37

Simple Reassurance..

- Instruct patients to breathe abdominally
 - Focus on diaphragm rather than chest wall
 - Slows rate
 - Distracts patient
 - Provides a sense of self control
- Abandon all if symptoms do not settle quickly



38